




**The Casualty Actuarial Society is committed to adhering strictly to the letter and spirit of the antitrust laws. Seminars conducted under the auspices of the CAS are designed solely to provide a forum for the expression of various points of view on topics described in the programs or agendas for such meetings.**

**Under no circumstances shall CAS seminars be used as a means for competing companies or firms to reach any understanding – expressed or implied – that restricts competition or in any way impairs the ability of members to exercise independent business judgment regarding matters affecting competition.**

**It is the responsibility of all seminar participants to be aware of antitrust regulations, to prevent any written or verbal discussions that appear to violate these laws, and to adhere in every respect to the CAS antitrust compliance policy.**



# BAYESIAN LOSS DEVELOPMENT: CREATING AN INFORMATIVE PRIOR

CAS Casualty Loss Reserve Seminar – September 19-20, 2016

Dave Clark  
Munich Reinsurance America, Inc

Munich RE  SM


# Agenda



1. Loss Development in Reinsurance Pricing
2. Bayesian Theory and Mathematics (optional)
3. Practical Implementation


## Cumulative Percent of Ultimate Reported

Examples of Fast and Slow Reporting



## Cumulative Percent of Ultimate Reported

Examples of Fast and Slow Reporting



# Loss Development Blending



Reinsurance pricing problem:

We have a loss development triangle from our client:

- May be sparse, not fully credible
- No tail beyond latest age in triangle

We have “benchmark” pattern from other sources:

- ISO / RAA / Reserving / Peer Companies
- Uncertain estimation and relevance for this client

# Loss Development Blending

(numbers for illustration only)



	Single Benchmark Example							
	12	24	36	48	60	72	84	96
1990	73	262	469	528	536	591	604	606
1991	148	346	391	502	522	514	567	
1992	99	198	219	394	408	430		
1993	118	255	352	412	581			
1994	275	415	645	803				
1995	261	446	637					
1996	130	471						
1997	148							
	<u>12-24</u>	<u>24-36</u>	<u>36-48</u>	<u>48-60</u>	<u>60-72</u>	<u>72-84</u>	<u>84-96</u>	<u>96-Ult</u>
1990	3.589	1.790	1.126	1.015	1.103	1.022	1.003	
1991	2.338	1.130	1.284	1.040	0.985	1.103		
1992	2.000	1.106	1.799	1.036	1.054			
1993	2.161	1.380	1.170	1.410				
1994	1.509	1.554	1.245					
1995	1.709	1.428						
1996	3.623							
Col. 1	1,104	1,922	2,076	1,836	1,466	1,105	604	
Col. 2	2,393	2,713	2,639	2,047	1,535	1,171	606	
Avg ATA	2.168	1.412	1.271	1.115	1.047	1.060	1.003	



Bayes' Theorem:

$$\pi(\theta|X) = \frac{f(X|\theta) \cdot \pi(\theta)}{\int f(X|\theta) \cdot \pi(\theta) d\theta}$$

This formula has three components:

$\pi(\theta)$  A distribution representing “prior” knowledge of the parameters  $\theta$

$f(X|\theta)$  A likelihood function representing the probability of observing the actual data  $X$  given a certain parameter set.

$\pi(\theta|X)$  The “posterior” probability of the parameters, revised based on the data





Tools for Evaluating the Mathematics:

- 1) Conjugate Families
- 2) Linear Approximation to Bayes Formula => Bühlmann-Straub
- 3) Numerical Approximation of the Formula
  - a) Quadrature integration (old method)
  - b) Simulation via MCMC (the new favorite)

Conjugate family has advantage of simple calculation and interpretability.



When the prior distribution  $\pi(\theta)$  and likelihood  $f(X|\theta)$  are chosen such that the posterior distribution  $\pi(\theta|X)$  has the same distribution form as the prior, then we have a *conjugate* relationship.

Common examples from the Exponential Family are:

$$\pi(\theta) \Rightarrow f(X|\theta)$$

Gamma  $\Rightarrow$  Poisson

Beta  $\Rightarrow$  Binomial

For the loss development pattern problem, we need a multivariate conjugate relationship.

Dirichlet  $\Rightarrow$  Multinomial

////////////////////////////////////

The Dirichlet distribution is a multivariate version of the beta distribution.

Instead of a yes/no probability of  $p$  or  $(1 - p)$ ,


we have a vector of probabilities  $\{p_1, p_2, p_3, \dots, p_k\}$ .

$$\pi(\mathbf{p}) = \frac{\Gamma(\alpha_1 + \alpha_2 + \dots + \alpha_k)}{\Gamma(\alpha_1) \cdot \Gamma(\alpha_2) \dots \Gamma(\alpha_k)} \cdot \prod_{j=1}^k p_j^{\alpha_j - 1} \cdot p_2^{\alpha_2 - 1} \dots p_k^{\alpha_k - 1}$$

$$E(p_i) = \frac{\alpha_i}{\sum \alpha_j} \quad 1 = \sum p_j$$

We can view this as a simulation of “k” gamma random variables, with a common scale parameter, which are then turned into percentages.

## Incremental Loss Development Pattern





The Dirichlet distribution gives us a prior pattern that matches the percents paid (or reported) in each incremental period. The increments are proportional to the  $\alpha_j$  parameters.

Given a new *observed* pattern from the client data, the “posterior” distribution simply adds a value to update the Dirichlet parameters.

$$\alpha_1, \alpha_2, \alpha_3, \dots, \alpha_k$$

becomes

$$(\alpha_1 + n_1), (\alpha_2 + n_2), (\alpha_3 + n_3), \dots, (\alpha_k + n_k)$$

However, this assumes we have a complete pattern from the client.

We do not; we have a series of incomplete patterns.



The Generalized Dirichlet distribution (Wong 1998) solves this challenge of incomplete data for us.

The Generalized form has twice as many parameters, but accommodates the incomplete data. Conveniently, this is also a conjugate form – meaning that the posterior distribution is again Generalized Dirichlet, with adjusted parameters.

The  $\alpha$ 's are incremental losses, the  $\beta$ 's are cumulative losses.

$$\{\alpha_1, \alpha_2, \alpha_3, \dots, \alpha_k, \beta_1, \beta_2, \beta_3, \dots, \beta_k\}$$

This has the remarkable interpretation that  $\left\{1 + \frac{\alpha_j}{\beta_j}\right\}_{j=1}^k$  is the sequence of age-to-age factors.



In addition to the uncertainty in the “prior” benchmark pattern, we need a measure of the process variance in the client triangle.

This is incorporated via a “dispersion” or variance/mean parameter  $\phi$ . Equivalent to the factor used in GLM or Bootstrapping for chain ladder.

$$\textit{Credibility } K = \frac{\textit{Expected Process Variance}}{\textit{Variance of Hypothetical Means}}$$

In the  $N/(N+K)$  formula, the “K” acts as ballast. It can be interpreted as counts or dollars depending upon the application.



Bayesian Combinations:

General Dirichlet:

$$ATA_d = \frac{\alpha_{k-d} + \beta_{k-d}}{\beta_{k-d}}$$

Chainladder:

$$ATA_d = \frac{\sum_{t=1}^{k-d} C_{t,d+1}}{\sum_{t=1}^{k-d} C_{t,d}}$$

Blended:

$$ATA_d = \frac{\phi \cdot (\alpha_{k-d} + \beta_{k-d}) + \sum_{t=1}^{k-d} C_{t,d+1}}{\phi \cdot \beta_{k-d} + \sum_{t=1}^{k-d} C_{t,d}}$$



# Credibility Blending Formula



The credibility blending becomes a simple dollar-weighted average.

If you can calculate an age-to-age factor, then you can do a Bayesian model!

		<u>Example of Blending Client and Benchmark Patterns</u>							
	<u>12-24</u>	<u>24-36</u>	<u>36-48</u>	<u>48-60</u>	<u>60-72</u>	<u>72-84</u>	<u>84-96</u>	<u>96-Ult</u>	
<u>ATA from Triangle</u>									
Col. 1	1,104	1,922	2,076	1,836	1,466	1,105	604	-	
Col. 2	2,393	2,713	2,639	2,047	1,535	1,171	606	-	
<b>ATA</b>	<b>2.168</b>	<b>1.412</b>	<b>1.271</b>	<b>1.115</b>	<b>1.047</b>	<b>1.060</b>	<b>1.003</b>		
<u>Benchmark Pattern</u>									
Col. 1	1,419	2,027	2,546	2,933	3,383	3,633	3,717	3,042	
Col. 2	4,000	4,000	4,000	4,000	4,000	4,000	4,000	4,000	
<b>ATA</b>	<b>2.819</b>	<b>1.973</b>	<b>1.571</b>	<b>1.364</b>	<b>1.182</b>	<b>1.101</b>	<b>1.076</b>	<b>1.315</b>	
<u>Blended Pattern</u>									
Col. 1	2,523	3,949	4,622	4,769	4,849	4,738	4,321	3,042	
Col. 2	6,393	6,713	6,639	6,047	5,535	5,171	4,606	4,000	
<b>ATA</b>	<b>2.534</b>	<b>1.700</b>	<b>1.436</b>	<b>1.268</b>	<b>1.141</b>	<b>1.091</b>	<b>1.066</b>	<b>1.315</b>	



“Conjugate priors... have the desirable feature that prior information can be viewed as ‘fictitious sample information’ in that it is combined with the sample in exactly the same way that additional sample information would be combined.

“The only difference is that the prior information is ‘observed’ in the mind of the researcher, not in the real world.”

- Bayesian Econometric Methods; Koop, Poirier & Tobias



How do we create the prior distribution  $\pi(\theta)$  ?

- 1) Empirical Bayes - estimate from collection of available patterns
- 2) Elicit ranges from expert users: e.g., select slow/medium/fast patterns
- 3) Reverse engineer – what prior is implied by the credibility percents that have been applied by users on actual accounts?

Different form for pricing and reserving actuaries?



## 1) Market Heterogeneity

“...the market experience is not fully relevant to a particular client. This is usually captured by the spread, or heterogeneity, of the client risk premiums around the standard market rate.”

## 2) Estimation Uncertainty


“...although the market rate is typically computed from a larger data set than that of a client, it, too, is based on a loss database of limited size and is therefore affected by the same type of uncertainty.”

- Parodi & Bonche

“Uncertainty-Based Credibility and its Applications” *Variance* 2010

# Estimate from collection of available patterns

## Products Liability



# Extending the Model



We can extend this model further by including mixtures of prior distributions.

Perhaps we know that companies are naturally grouped into Fast, Medium, or Slow payment patterns. But we do not know to which group our client belongs.

	<u>Cumulative Loss Development Factors</u>							
	<u>12</u>	<u>24</u>	<u>36</u>	<u>48</u>	<u>60</u>	<u>72</u>	<u>84</u>	<u>96</u>
Fast	14.014	4.930	2.607	1.759	1.406	1.263	1.191	1.155
Medium	21.950	7.787	3.946	2.512	1.842	1.558	1.415	1.315
Slow	49.240	15.860	7.407	4.163	2.706	2.057	1.750	1.567

# Extending the Model



We assign initial weights to each of the groupings (perhaps 33%/33%/33%) and then apply Bayes' theorem to update the weights.

<u>Bayesian Updating of Probabilities</u>						
	LogLikelihood	Difference in LL	Relative Likelihood	Original Weights	Revised Weights	
	A	$B=A-\text{Max}(A)$	$C=\exp(B)$	D	$E=C*D/\text{Avg}( C )$	
Slow	-4.61	-0.77	0.464	33.33%	20.41%	
Baseline	-4.06	-0.21	0.810	33.33%	35.61%	
Fast	-3.84	0.00	1.000	33.33%	43.98%	
			0.758	100.00%	100.00%	

This allows us to adjust our “tail” based on which group is closest to our client’s data.



- Credibility in Loss Development pattern selection has benefits
  - Stability in estimation – therefore can break data into small homogeneous pieces
  - Consistency in pricing
  - Even very sparse data from a client can update the benchmark
- The Bayesian framework can be implemented practically for pricing
- The Bayesian framework can be extended to include benchmarks for every uncertain parameter



## Selected References

---

Bolstad, William M., “Introduction to Bayesian Statistics,” Wiley, 2007.


Mildenhall Stephen J., “A Multivariate Bayesian Claim Count Development Model with Closed Form Posterior and Predictive Distributions,” *CAS Forum* Winter 2006, 451-493.

Wong, Tzu-Tsung, “Generalized Dirichlet Distribution in Bayesian Analysis,” *Applied Mathematics and Computation* 97 (1998), 165-181.

---

*Hot off the presses:*

Clark, David R., “Introduction to Bayesian Loss Development,” *CAS Forum* 2016.



Dave Clark

[daveclark@munichreamerica.com](mailto:daveclark@munichreamerica.com)

<https://twitter.com/DaveclarkR>

© Copyright 2016 Munich Reinsurance America, Inc. All rights reserved. "Munich Re" and the Munich Re logo are internationally protected registered trademarks. The material in this presentation is provided for your information only, and is not permitted to be further distributed without the express written permission of Munich Reinsurance America, Inc. or Munich Re. This material is not intended to be legal, underwriting, financial, or any other type of professional advice. Examples given are for illustrative purposes only. Each reader should consult an attorney and other appropriate advisors to determine the applicability of any particular contract language to the reader's specific circumstances.

Munich RE  SM